

## Hanson & Bussière 1998 - Table 4 & 5

**Table 4**  
**Correlations with recidivism for each category of predictor**

*Note*

Values represent average correlations  $\pm$  95% confidence interval.

Predictor	Type of recidivism		
	Sexual	Nonsexual violence	Any
Criminal lifestyle	.12 $\pm$ .02	.16 $\pm$ .03	.21 $\pm$ .02
Sexual deviance	.19 $\pm$ .02	.01 $\pm$ .03	.12 $\pm$ .02
Psychological maladjustment	.01 $\pm$ .03	.02 $\pm$ .08	.02 $\pm$ .03
Negative clinical presentation	.00 $\pm$ .07		.15 $\pm$ .07
Failure to complete treatment	.17 $\pm$ .07	.08 $\pm$ .09	.20 $\pm$ .07

**Table 5**  
**Combined risk scales**

*Note*

CI = confidence interval;  $n$  = total aggregated sample size.

<sup>a</sup> Values in parentheses represent number of studies.

\*\*  $p < .01$

Variable	<i>Mdn</i>	$r_+$	95% CI	<i>Q</i>		$n^a$	
<b>Sexual recidivism</b>							
Clinical assessment	.04	.10	.05 - .15	16.10		1,453	(10)
Statistical	.44	.46	.40 - .52	15.06	**	684	(6)
<b>Nonsexual violent recidivism</b>							
Clinical assessment	.06	.06	-.02 - .14	1.10		544	(3)
Statistical	.45	.46	.37 - .54	.65		343	(3)
<b>General recidivism</b>							
Clinical assessment	.11	.14	.08 - .20	17.81	**	1,067	(8)
Statistical	.42	.42	.36 - .51	.87		453	(5)